

Common

Abdominal pains and cramp
Agoraphobia
Anxiety
Breathing difficulties
Blurred vision
Changes in perception (faces distorting and inanimate objects moving)
Depression
Distended abdomen
Dizziness
Extreme lethargy
Fears
Feelings of unreality
Flu-like symptoms
Heavy limbs
Heart palpitations
Hypersensitivity to light
Indigestion
Insomnia
Irritability
Lack of concentration
Lack of co-ordination
Loss of balance
Loss of memory
Muscular aches and pains
Nausea
Nightmares
Panic attacks
Rapid mood changes
Restlessness
Severe headaches
Shaking
Seeing spots before the eyes
Sore eyes
Sweating
Tightness in the chest
Tightness in the head

Less Common

Aching jaw
Craving for sweet food
Constipation
Depersonalization (a feeling of not knowing who you are)
Diarrhea
Difficulty swallowing
Feeling of the ground moving
Hallucinations (auditory and visual)
Hyperactivity
Hypersensitivity to sound
Incontinence of urine, or frequency and urgency
Increased saliva
Loss or changes in appetite
Loss of taste or metallic taste
Menstrual problems (painful periods, irregular periods, cessation of periods)
Morbid thoughts
Numbness in any part of the body
Outbursts of rage or aggression
Paranoia
Painful scalp
Persistent, unpleasant memories
Pins and needles
Rapid changes in body temperature
Sexual problems (changes in libido)
Skin problems (dryness, itchiness, rashes, slow healing)
Sore mouth and tongue
Speech difficulties
Suicidal thoughts
Tinnitus (buzzing or ringing in the ears)
Unusually sensitive (unable to watch the news on television or read the newspaper)
Vaginal discharge
Uncharacteristic behavior such as shoplifting
Vomiting
Weight loss or gain

Rare

Blackouts -an episode where the person has no recall of their activity.(This is rare with low dose use, but less rare when large amounts have been taken)
Bleeding from the nose
Bleeding between menstrual cycles
Burning along the spine
Burning sensation around the mouth
Discharge from the breasts

Falling hair
Hemorrhoids
Hypersensitivity to touch
Rectal bleeding
Sinus pain
Seizures (fits) Almost unknown if people reduce gradually. More common for people using high doses who stop suddenly.
Sensitive or painful teeth
Swollen breasts